

The Generalized Heisenberg-Virasoro algebra

Dong Liu¹, Linsheng Zhu²

¹ Department of Mathematics, Huzhou Teachers College, Zhejiang Huzhou, 313000, China

² Department of Mathematics, Changshu Institute of Technology, Jiangsu Changshu, 215500, China

© Higher Education Press and Springer-Verlag 2009

Abstract In this paper, we mainly study the generalized Heisenberg-Virasoro algebra. Some structural properties of the Lie algebra are obtained.

Keywords The generalized Heisenberg-Virasoro algebra, central extension, automorphisms

MSC 17B56; 17B68

1 Introduction

The twisted Heisenberg-Virasoro algebra H_{Vir} has been first studied by Arbarello et al. in Ref. [1], where a connection is established between the second cohomology of certain moduli spaces of curves and the second cohomology of the Lie algebra of differential operators of order at most one:

$$L_{HV} = \{f(t)\frac{d}{dt} + g(t)|f, g \in \mathbb{C}[t, t^{-1}]\}.$$

As a vector space over $\mathbb{C}$, H_{Vir} has a basis $\{L(m), I(m), C_L, C_I, C_{LI}, m \in \mathbb{Z}\}$, subject to the following relations:

$$\begin{aligned} [L(m), L(n)] &= (n-m)L(m+n) + \delta_{m+n,0}\frac{1}{12}(m^3-m)C_L; \\ [I(m), I(n)] &= n\delta_{m+n,0}C_I; \\ [L(m), I(n)] &= nI(m+n) + \delta_{m+n,0}(m^2-m)C_{LI}; \\ [H_{Vir}, C_L] &= [H_{Vir}, C_I] = [H_{Vir}, C_{LI}] = 0. \end{aligned}$$

Clearly the Heisenberg algebra $H = \mathbb{C}\{I(m), C_I \mid m \in \mathbb{Z}\}$ and the Virasoro algebra $Vir = \mathbb{C}\{L(m), C_L \mid m \in \mathbb{Z}\}$ are subalgebras of H_{Vir} .

Arbarello et al. (in Ref. [1]) also proved that any irreducible highest weight module for H_{Vir} is isomorphic to the tensor product of an irreducible

* Received Oct. 10, 2008; accepted Feb. 25, 2009

noindent module for the Virasoro algebra and an irreducible module for the infinite-dimensional Heisenberg algebra when the central element of the Heisenberg subalgebra acts in a non-zero way. The structure of the irreducible representations for H_{Vir} at level zero was studied in Ref. [3]. The Harish-Chandra modules over H_{Vir} were classified in Refs. [11, 12]. Some structure properties and representations of the twisted Heisenberg-Virasoro Lie algebra were obtained in Refs. [7, 8, 9, etc.].

Recently, a number of new classes of infinite-dimensional simple Lie algebras over a field of characteristic 0 were discovered by several authors. Among those algebras, are the generalized Witt algebras, the generalized Virasoro algebras introduced in Ref. [16], which are one-dimensional universal central extensions of some generalized Witt algebras, and the Lie algebras of generalized Weyl type introduced and studied in Refs. [15, 17, etc.].

Motivated by the above algebras, we introduce a new Lie algebra, the generalized Heisenberg-Virasoro algebra (see Definition 1 in Section 2), which is a generalization of the twisted Heisenberg-Virasoro Lie algebra H_{Vir} from the integer ring $\mathbb{Z}$ to an additive subgroup M of a field $\mathbb{F}$ (The special case $M = \mathbb{Z}^2$ was studied in Ref. [18]). Its Verma modules were studied in Ref. [14]. In this paper, we mainly study some properties of this Lie algebra.

The paper is organized as follows. In Section 2, we recall some notions of generalized Witt algebras and the twisted Heisenberg-Virasoro algebra, and then introduce the definition of the generalized Heisenberg-Virasoro algebra. In Section 3, we prove that the generalized Heisenberg-Virasoro algebra is the universal extension of the Lie algebra of generalized differential operators on a circle of order at least one: $L = \mathbb{F}\{t^x\partial, t^y \mid x, y \in A, \partial \in T\}$. In Section 4 and Section 5, we determine derivations and automorphisms of the generalized Heisenberg-Virasoro algebra.

Throughout this paper, $\mathbb{F}$ denotes a field of characteristic zero, and $\mathbb{F}^* = \mathbb{F} \setminus \{0\}$. $\mathbb{C}$, $\mathbb{Z}$, $\mathbb{N}$, $\mathbb{Z}^+$ denote the field of complex numbers, the set of all integers, the set of all nonnegative integers, the set of all positive integers, respectively.

2 The generalized Heisenberg-Virasoro algebra

First we recall some notions of the generalized Witt algebras as defined in Refs. [4, 10].

Let $A (\neq 0)$ be an abelian group and T a vector space over $\mathbb{F}$. In this paper, we are only interested in the case of $\dim T = 1$, so we always assume that $T = \mathbb{F}\partial$ throughout this paper. We denote by $\mathbb{F}A$ the group algebra of A over $\mathbb{F}$. The elements $t^x, x \in A$, form a basis of the algebra, and the multiplication is defined by $t^x t^y = t^{x+y}$. We shall write 1 instead of t^0 . The

use the following notations:

$$\varphi(\partial, x) = \langle \partial, x \rangle = \partial(x)$$

for arbitrary $x \in A$.

There is a unique $\mathbb{F}$ -linear map $W \times W \rightarrow W$ sending $(t^x \partial, t^y \partial)$ to

$$[t^x \partial, t^y \partial] := t^{x+y} \partial(y - x) \partial \quad (2.1)$$

for arbitrary $x, y \in A$. This map makes $W = W(A, T, \varphi)$ into a Lie algebra which is called *generalized Witt algebra*. Setting $W_x = t^x T, x \in A$, then

$$W = \oplus_{x \in A} W_x$$

is a graded Lie algebra.

We say that φ is nondegenerate if $A_0 := \{x \in A : \partial(x) = 0\} = 0$.

The following theorem is due to N. Kawamoto (Ref. [10]).

Theorem 2.1 (Ref. [10]) *Suppose that characteristic of $\mathbb{F}$ is 0. Then $W = W(A, T, \varphi)$ is a simple Lie algebra if and only if φ is nondegenerate.*

In this paper, we always suppose that φ is nondegenerate.

In Ref. [5], the second cohomology group of the generalized Witt algebra was constructed.

Theorem 2.2 (Ref. [5]) *Let $W = W(A, T, \varphi)$ be a simple generalized Witt algebra and $T = \mathbb{F} \partial$ one-dimensional, then $H^2(W, \mathbb{F})$ is 1-dimensional and is spanned by the cohomology class $[\psi]$, where $\psi : W \times W \rightarrow \mathbb{F}$ is the 2-cocycle defined by*

$$\psi(t^x \partial, t^y \partial) = \delta_{x+y,0} \partial(x)^3, \quad x, y \in A. \quad (2.2)$$

The generalized Virasoro algebra is the universal central extension of the generalized Witt algebra when φ is nondegenerate (Ref. [16]).

By definition, the *generalized Virasoro algebra* Vir is a Lie algebra generated by $\{L(x) = t^x \partial, x \in A\}$ and C_L subject to the following relations:

$$\begin{aligned} [L(x), L(y)] &= \partial(y - x) L(x + y) + \delta_{x+y,0} \frac{1}{12} (\partial(x)^3 - \partial(x)) C_L \\ [L(x), C_L] &= 0. \end{aligned}$$

Now we introduce the algebra of generalized differential operators in Refs. [15, 16].

Let $\mathcal{D}$ be the associative algebra generated by the elements $\{t^x, x \in A\}$ and the element ∂ , subject to the following relation:

$$\partial t^x - t^x \partial = \partial(x) t^x, \quad \forall x \in A. \quad (2.3)$$

The algebra $\mathcal{D}$ is called the *algebra of generalized differential operators*. De-

basis $\{\partial\}$. Define the pairing $\varphi : T \times A \rightarrow \mathbb{F}$ by setting $\langle \partial, m \rangle = m$. Then φ is nondegenerate and the algebra of generalized differential operators is just the algebra of differential operators with coefficients in the Laurent polynomial ring.

As a vector space over $\mathbb{F}$, $\mathcal{D}$ has a basis $\{t^x \partial^m, x \in A, \partial \in T, m \in \mathbb{N}\}$. It is easy to see that the following relation holds in the associative algebra $\mathcal{D}$.

$$(t^x \partial^m)(t^y \partial^n) = t^{x+y} \sum_{i=0}^m \binom{m}{i} \partial(y)^i \partial^{m+n-i}, \quad \forall x, y \in A, m, n \in \mathbb{N}. \quad (2.5)$$

By (2.5) we can see that the generalized Witt algebra $W = W(A, T, \varphi)$ is a Lie subalgebra of $\mathcal{D}^-$, the Lie algebra of generalized differential operators.

The above results lead us to the following definition.

Definition 1. Let $T = \mathbb{F}\partial$ and $W = W(A, T, \varphi)$ be a simple generalized Witt algebra. The generalized Heisenberg-Virasoro algebra $\mathcal{L}$ is a Lie algebra generated by $\{L(x) = t^x \partial, I(x) = t^x, C_L, C_I, C_{LI}\}$, subject to the following relations:

$$\begin{aligned} [L(x), L(y)] &= \partial(y-x)L(x+y) + \delta_{x+y,0} \frac{1}{12} (\partial(x)^3 - \partial(x)) C_L; \\ [I(x), I(y)] &= \partial(y) \delta_{x+y,0} C_I; \\ [L(x), I(y)] &= \partial(y) I(x+y) + \delta_{x+y,0} (\partial(x)^2 - \partial(x)) C_L; \\ [\mathcal{L}, C_L] &= [\mathcal{L}, C_I] = [\mathcal{L}, C_{LI}] = 0. \end{aligned}$$

The Lie algebra $\mathcal{L}$ has a generalized Heisenberg subalgebra and a generalized Virasoro subalgebra intertwined with a 2-cocycle.

Moreover, we shall prove that the generalized Heisenberg-Virasoro algebra $\mathcal{L}$ is the universal central extension of the Lie algebra of generalized differential operators of order at least one: $\mathcal{D}_1 = \mathbb{F}\{t^x \partial, t^y \mid x, y \in A, \partial \in T\}$.

3 The universal central extension of the generalized Heisenberg-Virasoro algebra

Now we consider the Lie subalgebra $\mathcal{D}_1 = \mathbb{F}\{t^x \partial, t^y \mid x, y \in A\}$ of $\mathcal{D}^-$, the Lie algebra of generalized differential operators.

Theorem 3.1 $\dim H^2(\mathcal{D}_1, \mathbb{F}) = 3$.

Proof. Let α be any 2-cocycle of $\mathcal{D}_1$ and we set $\alpha_{x,y} = \alpha(t^x \partial, t^y \partial)$, $\alpha_x^y = \alpha(t^x \partial, t^y)$ and $\alpha^{x,y} = \alpha(t^x, t^y)$. Considering $\alpha([t^z \partial, t^x], t^y)$, $\alpha([t^z \partial, t^x \partial], t^y)$, $\alpha([t^z \partial, t^x \partial], t^y \partial)$ and using the relations:

$$[t^x \partial, t^y \partial] = \partial(y-x)t^{x+y} \partial, \quad [t^x \partial, t^y] = \partial(y)t^{x+y}, \quad [t^x, t^y] = 0,$$

and the cocycle rule, we get the following

$$\partial(x-z)\alpha_{z+x,y} + \partial(y-x)\alpha_{x+y,z} + \partial(z-y)\alpha_{y+z,x} = 0. \quad (3.3)$$

The relation (3.1) immediately implies

$$\alpha^{x,y} = \frac{\partial(y)}{\partial(x_0)} \delta_{x+y,0} \alpha^{-x_0,x_0}, \forall x, y \in A, \quad (3.4)$$

for some $x_0 \in A$ such that $\partial(x_0) \neq 0$.

By (3.4), $\alpha^{x,y}$ is determined by nontrivial 2-cocycle $\psi_1 : \mathcal{D}_1 \times \mathcal{D}_1 \rightarrow \mathbb{F}$ defined by $\psi_1(t^x, t^y) = \delta_{x+y,0} \partial(y)$, for all $x, y \in A$.

From the proof of Theorem 6.1 in Ref. [5], we see that, up to a coboundary, it follows that $\alpha_{x,y}$ is determined by nontrivial 2-cocycle ψ_2 , where $\psi_2(t^x \partial, t^y \partial) = \delta_{x+y,0} (\partial(x)^3 - \partial(x))$, for all $x, y \in A$.

Setting $y = 0$ in (3.2) we deduce that

$$\alpha_x^0 = 0, \quad \forall x \in A. \quad (3.5)$$

Let $g : \mathcal{D}_1 \rightarrow \mathbb{F}$ be the linear function $g(t^x \partial) = 0$ and $g(t^x) = \frac{1}{\partial(x)} \alpha(\partial, t^x)$ if $x \neq 0$ and $g(t^0) = 0$. Then g induces a 2-coboundary $\psi_g(u, v) = g([u, v])$, $u, v \in \mathcal{D}_1$. Hence by replacing α with the 2-coboundary $\alpha - \psi_g$, we may assume that $\alpha(\partial, t^y) = 0$ for all $y \neq 0$. Hence by (3.5), we obtain

$$\alpha_0^x = 0, \quad \forall x \in A.$$

Setting $z = 0$ in (3.2), we have

$$\alpha_x^y = 0, \quad \forall x, y \in A \text{ and } x + y \neq 0. \quad (3.6)$$

On the other hand, setting $y = -z - x$ in (3.2), we have

$$\partial(x-z)\alpha_{x+z}^{-x-z} = \partial(x+z)(\alpha_x^{-x} - \alpha_z^{-z}), \quad \forall x, z \in A. \quad (3.7)$$

Let $x \in A$ be such that $\partial(x) \neq 0$. By substituting kx for x and lx for z , we obtain

$$(k-l)f(k+l) = (k+l)(f(k) - f(l)), \quad (3.8)$$

where $f(k) = \alpha_{kx}^{-kx}$, $k \in \mathbb{Z}$.

It follows from (3.8) that all the values $f(k)$ can be computed if $f(1)$ and $f(2)$ are known. Hence the general solution (3.8) is given by $f(k) = ak + bk^2$ for some constants $a, b \in \mathbb{F}$.

Setting $k = 1$ and $k = 2$, we have

$$\alpha_{kx}^{-kx} = \frac{k}{2}(4\alpha_x^{-x} - \alpha_{2x}^{-2x}) + \frac{k^2}{2}(\alpha_{2x}^{-2x} - 2\alpha_x^{-x}) \quad (3.9)$$

for all $k \in \mathbb{Z}$, provided that $\partial(x) \neq 0$.

Let x_0 be a nonzero element of A . Using similar considerations in Page

Then α_x^y is determined by a nontrivial 2-cocycle $\psi_3 : \mathcal{D}_1 \times \mathcal{D}_1 \rightarrow \mathbb{F}$ defined by $\psi_3(t^x \partial, t^y) = \delta_{x+y,0} \partial(x)^2$ and the 2-coboundary $\psi'_3(t^x \partial, t^y) = \delta_{x+y,0} \partial(x)$ for all $x, y \in A$. $\blacksquare$

Corollary 3.2 *The generalized Heisenberg-Virasoro algebra $\mathcal{L}$ is the universal central extension of the Lie algebra $\mathcal{D}_1$ of generalized differential operators of order at least one.* $\blacksquare$

4 Derivations of the generalized Heisenberg-Virasoro algebra

In this section we shall determine all derivations of the generalized Heisenberg-Virasoro algebra $\mathcal{L}$.

First, we recall a result about derivations in Ref. [2].

Proposition 4.1 (Ref. [2]) *If L is a perfect Lie algebra and $\hat{L}$ is a universal central extension of L , then every derivation of L lifts to a derivation of $\hat{L}$. If L is centerless, the lift is unique and $\text{Der}(\hat{L}) = \text{Der}(L)$.*

Due to Corollary 3.2 in Section 3 and Proposition 4.1, we shall just determine all derivations of $\mathcal{D}_1 = \mathbb{F}\{t^x \partial, t^y \mid x, y \in A, \partial \in T\}$, the Lie subalgebra of $\mathcal{D}^-$.

We now describe two kinds of derivations of degree 0, which are outer derivations of $\mathcal{D}_1$.

The linear maps $\sigma_1, \sigma_2, \sigma_3 : \mathcal{D}_1 \rightarrow \mathcal{D}_1$ defined by

$$\sigma_1(L(x)) = \partial(x)I(x), \quad \sigma_1(I(x)) = 0, \quad \forall x \in A,$$

and

$$\sigma_2(L(x)) = I(x), \quad \sigma_2(I(x)) = 0, \quad \forall x \in A,$$

and

$$\sigma_3(L(x)) = 0, \quad \sigma_3(I(x)) = I(x), \quad \forall x \in A,$$

are outer derivations of $\mathcal{D}_1$.

Let $\mu : A \rightarrow \mathbb{F}$ be an additive map, then the linear map $\xi_\mu : \mathcal{D}_1 \rightarrow \mathcal{D}_1$ defined by

$$\xi_\mu(L(x)) = \mu(x)L(x), \quad \xi_\mu(I(x)) = \mu(x)I(x), \quad \forall x \in A,$$

is also a derivation of degree 0. It is clear that it is an outer derivation if $\mu \neq k\partial$ for any $k \in \mathbb{F}$. In fact, if $\mu = k\partial$ for some $k \in \mathbb{F}$, then $\xi_\mu = \text{ad}(k\partial)$.

Let A be an abelian group, $\mathfrak{g}$ an A -graded Lie algebra, and V an A -graded left $\mathfrak{g}$ -module. We also denote by L_x resp. V_x the homogeneous components of $\mathcal{D}_1$ resp. V for any $x \in A$.

Proposition 4.2 (Refs. [5, 6]) *Every derivation $D \in \text{Der}(\mathfrak{g}, V)$ can be written as*

$$D = \sum_{x \in A} D_x, \quad D_x \in \text{Der}(\mathfrak{g}, V)_x,$$

where $\text{Der}(\mathfrak{g}, V)_x = \{\sigma \in \text{Der}(\mathfrak{g}, V) \mid \sigma(L_y) \subset V_{x+y}, \forall y \in A\}$.

A derivation $D : \mathfrak{g} \rightarrow V$ is called *locally inner* (Ref. [5]) if it is a sum (may be infinite sum) of inner derivations.

Proposition 4.3 (Refs. [5, 6]) *Suppose that the following conditions hold.*

- (1) $H^1(\mathfrak{g}_0, V_x) = 0$ for $x \neq 0$;
- (2) $\text{Hom}_{\mathfrak{g}_0}(L_x, V_y) = 0$ for $x \neq y$.

Then $\text{Der}(\mathfrak{g}, V)_x$, $x \neq 0$, consists of inner derivations and consequently

$$\text{Der}(\mathfrak{g}, V) = \text{Der}(\mathfrak{g}, V)_0 + \text{Der}'(\mathfrak{g}, V),$$

where $\text{Der}'(\mathfrak{g}, V)$ is the space of locally inner derivations $\mathfrak{g} \rightarrow V$.

We now use Proposition 4.3 to prove the following result.

Proposition 4.4

$$\text{Der}(\mathcal{D}_1) = \text{ad}(\mathcal{D}_1) + \text{Der}(\mathcal{D}_1)_0.$$

Proof. By direct calculation, see Ref. [5]. ■

Theorem 4.5 *Assume that $A \neq 0$ and φ is nondegenerate. Then*

$$\text{Der}(\mathcal{D}_1)_0 = \text{ad}(\mathcal{D}_1)_0 + \sum_{\mu \in \text{Hom}(A, \mathbb{F})} \mathbb{F}\xi_\mu + \sum_{i=1}^3 \mathbb{F}\sigma_i,$$

Proof. Since φ is nondegenerate, we can suppose that $\partial(x) \neq 0$ if $x \neq 0$.

For any $D \in \text{Der}(\mathcal{D}_1)_0$ and $x \in A$, we suppose

$$D(t^x) = t^x(\alpha_x \partial + \beta_x), \alpha_x, \beta_x \in \mathbb{F}. \quad (4.1)$$

For any $y \in A$, by the fact that $[t^x, t^y] = 0$, we obtain that

$$\alpha_x \partial(y) - \partial(x) \alpha_y = 0. \quad (4.2)$$

Choose $x_0 \in A$ such that $\partial(x_0) = 1$ and set $a_0 = \alpha_{x_0}$. So by (4.2) we have

$$\alpha_y = \partial(y) a_0, \quad (4.3)$$

and (4.1) becomes

$$D(t^x) = t^x(a_0 \partial(x) \partial + \beta_x), \quad a_0, \beta_x \in \mathbb{F}. \quad (4.4)$$

Now we suppose that

$$D(t^y \partial) = t^y(\gamma_y \partial + \lambda_y), \quad \gamma_y, \lambda_y \in \mathbb{F}. \quad (4.5)$$

Applying D to $[t^y \partial, t^x] = \partial(x) t^{x+y}$, we obtain

$$[D(t^y \partial), t^x] + [t^y \partial, D(t^x)] = \partial(x) D(t^{x+y}).$$

So $\partial(x)\partial(x-y)a_0 = \partial(x)\partial(x+y)a_0$ and $\gamma_y\partial(x) + \beta_x\partial(x) = \partial(x)\beta_{x+y}$ for all $x, y \in A$. Hence

$$a_0 = 0 \quad (4.6)$$

and

$$\beta_{x+y} = \beta_x + \gamma_y, \quad \forall x \neq 0, y \in A. \quad (4.7)$$

Substitute D by $D - \beta_0\sigma_3$, we can suppose $\beta_0 = 0$. Setting $y = -x$ in (4.7), we have

$$\beta_x = -\gamma_{-x}, \quad \forall x \in A. \quad (4.8)$$

So (4.4) and (4.5) become

$$D(t^x) = \beta_x t^x. \quad (4.9)$$

$$D(t^x\partial) = t^x(\gamma_x\partial + \lambda_x), \quad \beta_x = -\gamma_{-x}, \lambda_x \in \mathbb{F}. \quad (4.10)$$

Applying D to $[t^x\partial, t^y\partial] = \partial(y-x)t^{x+y}\partial$, we obtain

$$\gamma_{x+y} = \gamma_x + \gamma_y, \quad \text{and} \quad \partial(y-x)\lambda_{x+y} = \partial(y)\lambda_y - \partial(x)\lambda_x. \quad (4.11)$$

By (4.11) and (4.8) we have $\gamma_x = \beta_x$ since $\gamma_0 = 0$. Then $\beta : A \rightarrow \mathbb{F}$ defined by $\beta(x) = \beta_x$ is an additive map. Moreover by using the method as in Section 3 (see (3.7)-(3.9)) we can deduce that $\lambda_x = a\partial(x) + b$ for some $a, b \in \mathbb{F}$ from (4.11). Replacing D by $D - \xi_\beta - a\sigma_1 - b\sigma_2$, we infer that $D = 0$. ■

5 The automorphism group of $\mathcal{L}$

Now, we recall a result about automorphisms in Ref. [13].

Proposition 5.1 (Ref. [13]) *Let L be a perfect Lie algebra and $\hat{L}$ be its universal central extension. Every automorphism θ of L admits a unique extension $\hat{\theta}$ to an automorphism $\hat{L}$. Furthermore, the map $\theta \rightarrow \hat{\theta}$ is a group monomorphism.*

Due to Corollary 3.2 in Section 3 and Proposition 5.1, we shall just determine the automorphism group of $\mathcal{D}_1 = \mathbb{F}\{t^x\partial, t^y \mid x, y \in A, \partial \in T\}$. Set $I = \mathbb{F}\{t^x \mid x \in A\}$, then I is the unique maximal solvable ideal of $\mathcal{D}_1$. Clearly, if π is an automorphism of $\mathcal{D}_1$, then

$$\pi(I) = I. \quad (5.1)$$

Denote by $\mathfrak{r}$ the set of all inner automorphisms of $\mathcal{D}_1$, then $\mathfrak{r}$ is a normal subgroup of $\text{Aut}(\mathcal{D}_1)$ and $\mathfrak{r}$ is generated by $\exp(k \text{ad } t^x)$, $x \in A, k \in \mathbb{F}$.

For convenience, denote by $X(A)$ the group of characters of A , i.e., the group homomorphisms $A \rightarrow \mathbb{F}^*$. Set $\mathcal{L} = \{ \xi \in \mathbb{F}^* \mid \text{Span} \pi t^{\varepsilon x} \mid x \in A \} =$

Lemma 5.2 *Let $\pi \in \text{Aut}(\mathcal{D}_1)$, then there exists $\eta \in \mathfrak{r}$, such that*

$$\eta^{-1}\pi(\partial) = \varepsilon^{-1}\partial + a,$$

for some $\varepsilon \in \mathcal{E}$ and $a \in \mathbb{F}$.

Proof. Assume that

$$\pi(\partial) = \sum_{z \in A_1} \lambda_z t^z \partial + \sum_{w \in A_2} \gamma_w t^w,$$

where $\lambda_z, \gamma_w \in \mathbb{F}$ and A_1, A_2 are finite subsets of A . By (5.1), we can assume

$$\pi(t^x) = \sum_{y \in A_3} \nu(y) t^y,$$

where A_3 is a finite subset of A . Since $[\partial, t^x] = \partial(x)t^x$, we have

$$\left[\sum_{z \in A_1} \lambda_z t^z \partial + \sum_{w \in A_2} \gamma_w t^w, \sum_{y \in A_3} \nu(y) t^y \right] = \partial(x) \sum_{y \in A_3} \nu(y) t^y.$$

Then

$$\left[\sum_{z \in A_1} \lambda_z t^z \partial, \sum_{y \in A_3} \nu(y) t^y \right] = \partial(x) \sum_{y \in A_3} \nu(y) t^y. \quad (5.2)$$

So if $z \neq 0$, then $\lambda_z = 0$. Moreover, (5.2) becomes

$$[\lambda_0 \partial, \sum_{y \in A_3} \nu(y) t^y] = \partial(x) \sum_{y \in A_3} \nu(y) t^y. \quad (5.3)$$

So $\lambda_0 \partial(y) = \partial(x)$ for all $y \in A_3$. Since ∂ is nondegenerate, $x = \lambda_0 y$.

Hence

$$\pi(\partial) = \lambda_0 \partial + \sum_{z \in A_2} \gamma_z t^z \quad (5.4)$$

and

$$\pi(t^x) = \nu(\lambda_0^{-1}x) t^{\lambda_0^{-1}x}.$$

So $\pi(I) \subseteq I' = \text{Span}_{\mathbb{F}}\{t^{\lambda_0^{-1}x}, x \in A\}$. But $I = I'$, we deduce that $\varepsilon = \lambda_0^{-1} \in \mathcal{E}$.

Therefore

$$\pi(t^x) = \nu(\varepsilon x) t^{\varepsilon x}$$

and

$$\pi(\partial) = \varepsilon^{-1} \partial + \sum_{w \in A_2} \gamma_w t^w$$

where $\varepsilon \in \mathcal{E}$.

Let

Let $\chi \in X(A)$, $\varepsilon \in \mathcal{E}$, $a, b \in \mathbb{F}$, $c \in \mathbb{F}^*$, then there is a unique linear map

$$\theta = \theta(\chi, \varepsilon, a, b, c) : \mathcal{D}_1 \rightarrow \mathcal{D}_1$$

such that

$$\theta(t^x \partial) = \varepsilon^{-1} \chi(x) t^{\varepsilon x} \partial + (b \partial(x) + a) \chi(x) t^{\varepsilon x}, \quad \theta(t^y) = c \chi(y) t^{\varepsilon y} \quad (5.7)$$

for all $x \in A$.

It is straightforward to verify that θ is an automorphism of Lie algebra $\mathcal{D}_1$.

Lemma 5.3 *Let π be an automorphism of $\mathcal{D}_1$, then there exist $b \in \mathbb{F}$, $c \in \mathbb{F}^*$, $\chi \in X(A)$, $\varepsilon \in \mathcal{E}$, such that $\pi = \theta(\chi, \varepsilon, a, b, c)$.*

Proof. By Lemma 5.2, we can suppose that $\pi(\partial) = \varepsilon^{-1} \partial + a$ and $\pi(t^x) = \nu(x) t^{\varepsilon x}$ for some $a \in \mathbb{F}$, $\varepsilon \in \mathcal{E}$ and $\nu(x) \in \mathbb{F}$. By the fact that $[t^x \partial, t^y] = \partial(y) t^{x+y}$ we can deduce that $\pi(t^x \partial) \in L_{\varepsilon x}$. So

$$\pi(L_x) = L_{\varepsilon x}.$$

For each $x \in A$, we can suppose that

$$\pi(t^x \partial) = \varepsilon^{-1} \chi(x) t^{\varepsilon x} \partial + \tau(x) t^{\varepsilon x}, \quad (5.8)$$

$$\pi(t^x) = \lambda(x) t^{\varepsilon x}, \quad (5.9)$$

where $\chi(x), \lambda(x) \in \mathbb{F}^*$, $\tau(x) \in \mathbb{F}$. Moreover, $\chi(0) = 1$ and $\tau(0) = a$ since $\pi(\partial) = \varepsilon^{-1} \partial + a$. We claim that $\chi(x+y) = \chi(x)\chi(y)$ for all $x, y \in A$. It suffices to prove this for $x \neq y$. Then by applying π to

$$[t^x \partial, t^y \partial] = \partial(y-x) t^{x+y} \partial \quad (5.10),$$

we obtain

$$\begin{aligned} & [\varepsilon^{-1} \chi(x) t^{\varepsilon x} \partial + \tau(x) t^{\varepsilon x}, \varepsilon^{-1} \chi(y) t^{\varepsilon y} \partial + \tau(y) t^{\varepsilon y}] \\ &= \partial(y-x) (\varepsilon^{-1} \chi(x+y) t^{\varepsilon(x+y)} \partial + \tau(x+y) t^{\varepsilon(x+y)}). \end{aligned} \quad (5.11)$$

Therefore

$$\chi(x+y) = \chi(x)\chi(y) \quad (5.12)$$

and

$$\partial(y-x) \tau(x+y) = \partial(y) \tau(y) \chi(x) - \partial(x) \tau(x) \chi(y). \quad (5.13)$$

Suppose that $\tau(x) = f(x) \chi(x)$, then (5.13) becomes

$$\partial(y-x) f(x+y) = \partial(y) f(y) - \partial(x) f(x). \quad (5.14)$$

Fix $0 \neq x_0 \in \mathbb{F}$. By substituting x_0 and $2x_0$ for y in (5.14) respectively, we obtain

By substituting x_0 for x and $x + x_0$ for y in (5.14), we obtain

$$\partial(x)f(x + 2x_0) = (\partial(x) + \partial(x_0))f(x + x_0) - \partial(x_0)f(x_0). \quad (5.17)$$

Since φ is non-degenerate, by (5.15)-(5.17), we deduce that

$$f(x) = \frac{f(2x_0) - f(x_0)}{\partial(x_0)}\partial(x) + 2f(x_0) - f(2x_0).$$

So we can assume that $f(x) = b\partial(x) + a$ for some $b \in \mathbb{F}$.

By applying π to

$$[t^x\partial, t^y] = \partial(y)t^{x+y},$$

we obtain

$$[\varepsilon^{-1}\chi(x)t^{\varepsilon x}\partial + \tau(x)t^{\varepsilon x}, \lambda(y)t^{\varepsilon y}] = \partial(y)\lambda(x+y)t^{\varepsilon(x+y)}.$$

So

$$\chi(x)\lambda(y)\partial(y)t^{\varepsilon(x+y)} = \partial(y)\lambda(x+y)t^{\varepsilon(x+y)}.$$

Therefore

$$\lambda(x+y)\partial(y) = \chi(x)\lambda(y)\partial(y). \quad (5.18)$$

By setting $x = -y$ in (5.18), we obtain $\lambda(y) = c\chi(y)$ if $y \neq 0$, where $c = \lambda(0)$. Therefore $\lambda(x) = c\chi(x)$ for all $x \in A$ since $\chi(0) = 1$. $\blacksquare$

By Lemma 5.2 and Lemma 5.3, we have the following result.

Theorem 5.4 *Set*

$$\mathbf{aut}(\mathcal{D}_1) = \{\theta(\chi, \varepsilon, a, b, c) \mid \chi \in X(A), \varepsilon \in \mathcal{E}, a, b \in \mathbb{F}, c \in \mathbb{F}^*\},$$

then

$$Aut(\mathcal{D}_1) \cong \mathfrak{r} \rtimes \mathbf{aut}(\mathcal{D}_1).$$

Let $\theta(\chi_i, \varepsilon_i, a_i, b_i, c_i)$, $i = 1, 2$ be two automorphisms of $\mathcal{D}_1$ defined as in (5.7), then

$$\begin{aligned} & \theta(\chi_1, \varepsilon_1, a_1, b_1, c_1)\theta(\chi_2, \varepsilon_2, a_2, b_2, c_2) \\ &= \theta((\chi_1 \circ \varepsilon_2)\chi_2, \varepsilon_1\varepsilon_2, \varepsilon_2^{-1}a_1 + c_1a_2, b_1 + c_1b_2, c_1c_2). \end{aligned} \quad (5.19)$$

Moreover

$$\theta(\chi, \varepsilon, a, b, c)^{-1} = \theta(\chi^{-1} \circ \varepsilon^{-1}, \varepsilon^{-1}, -\varepsilon ac^{-1}, -bc^{-1}, c^{-1}). \quad (5.20)$$

The map $X(A) \rightarrow \mathbf{aut}(\mathcal{D}_1)$ defined by

Set

$$\mathfrak{a} = \{\theta(1, 1, a, b, 1) \mid a, b \in \mathbb{F}\}, \quad \mathfrak{c} = \{\theta(1, 1, 0, 0, c) \mid c \in \mathbb{F}^*\}.$$

Clearly

$$\mathfrak{a} \cong \mathbb{F} \times \mathbb{F}, \quad \mathfrak{c} \cong \mathbb{F}^*,$$

and $\mathfrak{a}, \mathfrak{c}$ are subgroups of $\mathbf{aut}(\mathcal{D}_1)$. Moreover $N\mathfrak{ac}$ is a normal group of $\mathbf{aut}(\mathcal{D}_1)$.

Denote by $\mathcal{A}$ the image of the endomorphism of $\mathbf{aut}(\mathcal{D}_1)$ defined by $\theta(\chi, \varepsilon, a, b, c) \mapsto \theta(1, \varepsilon, 0, 0, 1)$. Its kernel is $N\mathfrak{ac}$ and its restriction to $\mathcal{A}$ is the identity map. Hence $\mathbf{aut}(\mathcal{D}_1) \cong (N\mathfrak{ac}) \rtimes \mathcal{A}$ and $\mathcal{A} \cong \mathcal{E}$. Therefore we have the following result.

Theorem 5.5

$$\mathbf{aut}(\mathcal{D}_1) \cong X(A)\mathfrak{ac} \rtimes \mathcal{E}.$$

Acknowledgements Project is supported by the NNSF (Grant 10671027, 10701019), the ZJNSF(Grant Y607136, D7080080), the "Qianjiang Excellence Project" (No. 2007R10031) and the "New Century 151 Talent Project" (2008) of Zhejiang Province. Authors are grateful to the referee for correction in some errors and invaluable suggestions.

References

1. Arbarello E, DeConcini C, Kac V G, Procesi C. Moduli spaces of curves and representation theory, *Commun Math Phys*, 1988, 117: 1-36
2. Benkart G M, Moody R V. Derivations, central extensions and affine Lie algebras, *Algebras Groups Geom*, 1986, 3(4): 456-492
3. Billig Y. Representations of the twisted Heisenberg-Virasoro algebra at level zero, *Canad Math Bull*, 2003, 46(4): 529-537
4. Doković D Ž, Zhao Kaiming. Generalized Cartan type W Lie algebras in characteristic zero, *J Algebra*, 1997, 195: 170-210
5. Doković, D Ž, Zhao Kaiming. Derivations, isomorphisms, and second cohomology of generalized Witt algebras, *Trans Amer Math Soc*, 1998, 350(2): 643-664
6. Farnsteiner R. Derivations and extensions of finitely generated graded Lie algebras, *J. Algebra*, 1988, 118(1): 34-45
7. Fabbri M A, Moody R V. Irreducible representations of Virasoro-toroidal Lie algebras, *Comm Math Phys*, 1994, 159(1): 1-13
8. Fabbri M A, Okoh F. Representations of Virasoro-Heisenberg algebras and Virasoro-toroidal algebras, *Canad J Math*, 1999, 51(3): 523-545
9. Jiang Qifen, Jiang Cuipo. Representations of the twisted Heisenberg-Virasoro algebra and the full Toroidal Lie Algebras, *Algebra Colloquium*, 2007, 14(1): 117-134

12. Lv Rencai, Zhao Kaiming. Classification of irreducible weight modules over the twisted Heisenberg-Virasoro algebra, math-st/0510194
13. Pianzola A. Automorphisms of toroidal Lie algebras and their central quotients, J. Algebra Appl, 2002, 1(1): 113–121
14. Shen Ran, Jiang Qifen, Su Yucai. Verma Modules Over the Generalized Heisenberg-Virasoro Algebra, Comm. Alg, 2008, 38(4): 1464-1473.
15. Su Yucai, Zhao Kaiming. Simple algebras of Weyl type, Science in China A, 2001, 44: 419-426.
16. Su Yucai, Zhao Kaiming. Generalized Virasoro and super-Virasoro algebras and modules of the intermediate series, J Alg, 2002, 252: 1–19
17. Su Yucai, Zhao Kaiming. Structure of Lie algebras of Weyl type, Comm Alg, 2004, 32: 1052–1059
18. Xue Min, Lin Weiqiang, Tan Shaobin. Central extension, derivations and automorphism group for Lie algebras arising from the 2-dimensional torus, Journal of Lie Theory, 2005, 16: 139-153